

**PCT**WORLD INTELLECTUAL P  
Internation

INTERNATIONAL APPLICATION PUBLISHED UN

(51) International Patent Classification<sup>6</sup> :**C12N 15/49, C07K 14/16, C12N 15/62,  
15/57, A61K 39/21, C12N 15/86, 5/10 //  
9/52, 9/62, 9/64, 7/01****A1**

(1

**WO 9607741A1****1**

(43) International Publication Date: 14 March 1996 (14.03.96)

(21) International Application Number: PCT/CA95/00510

(22) International Filing Date: 7 September 1995 (07.09.95)

(30) Priority Data:  
08/301,915 7 September 1994 (07.09.94) US(71) Applicant: UNIVERSITE DE MONTREAL [CA/CA]; 2900  
Edouard-Montpetit, Montréal, Québec H3T 1J4 (CA).(72) Inventors: COHEN, Eric, A.; 4815 Wilson, Montréal, Québec  
H3X 3P3 (CA). BERGERON, Dominique; 3639 Laval,  
Montréal, Québec H2X 3E1 (CA). CHECROUNE, Florent;  
Apartment 202, 1101 Saint-Urbain, Montréal, Québec H2Z  
1Y6 (CA). YAO, Xiao-Jian; 3860 Barclay, Montréal,  
Québec H3S 1K7 (CA). PIGNAC-KOBINGER, Gary;  
2194A Mont-Royal, Montréal, Québec H2H 1K3 (CA).(74) Agent: COTE, France; Swabey Ogilvy Renault, Suite 1600,  
1981 McGill College Avenue, Montréal, Québec H3A 2Y3  
(CA).(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,  
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE,  
KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK,  
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT,  
BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL,  
PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,  
ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD,  
SZ, UG).**Published***With international search report.*

(54) Title: PROTEIN TARGETING INTO HIV VIRIONS BASED ON HIV-1 VPR FUSION MOLECULES

(57) Abstract

The present invention relates to a Vpr protein, a Vpx protein or fragments thereof which permit the development of chimeric molecules that can be specifically targeted into the mature HIV-1 and HIV-2 virions to affect their structural organization and/or functional integrity, thereby resulting in gene therapy for HIV-1 and HIV-2 infections. The present invention also relates to Vpr/Vpx protein fragments, p6 protein, p6 protein fragment, or functional derivatives thereof which interfere with the native Vpr/Vpx incorporation into HIV-1 and HIV-2 virions. The present invention also relates to treatment of HIV-1 and HIV-2 infections based on the proteins of the present invention.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/07233

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9101375	07-02-91	AU-B- 627063	13-08-92
		AU-B- 6042390	22-02-91
		EP-A- 0482053	29-04-92
		JP-T- 4506602	19-11-92
		US-A- 5365015	15-11-94
-----			
WO-A-9204456	19-03-92	AU-B- 657276	09-03-95
		AU-B- 8511491	30-03-92
		CA-A- 2091243	11-03-92
		EP-A- 0548164	30-06-93
		JP-T- 6502759	31-03-94
-----			
WO-A-9408449	28-04-94	AU-B- 5442894	09-05-94
-----			
WO-A-9212249	23-07-92	AU-B- 9113791	17-08-92
		BR-A- 9107191	14-06-94
		CA-A- 2096637	27-06-92
		EP-A- 0564524	13-10-93
		JP-T- 6504668	02-06-94
-----			
WO-A-9211371	09-07-92	AU-B- 660637	06-07-95
		AU-B- 9105691	22-07-92
		AU-B- 660638	06-07-95
		AU-B- 9106191	22-07-92
		EP-A- 0563157	06-10-93
		EP-A- 0563161	06-10-93
		WO-A- 9211372	09-07-92
		JP-T- 6503953	12-05-94
		JP-T- 6503954	12-05-94
		US-A- 5449764	12-09-95
-----			

- 1 -

PROTEIN TARGETING INTO HIV VIRIONS BASED ON  
HIV-1 VPR FUSION MOLECULES

TECHNICAL FIELD

This invention relates to two different  
5 approaches using the Vpr/Vpx protein or p6 protein for  
treatment of HIV-1 and/or HIV-2 infections.

BACKGROUND ART

Acquired Immune Deficiency Syndrome (AIDS) is a  
10 slow degenerative disease of the immune and nervous  
systems caused by the Human Immunodeficiency Virus  
(HIV). The life cycle of HIV lies at the heart of the  
AIDS pandemic. The spread of the disease is primarily  
determined by the infectious properties of this virus.  
15 Progressive lethal degeneration of the immune and  
central nervous systems results from long term chronic  
replication of this virus.

HIV belongs to a unique virus family, the  
retroviridae, a group of small, enveloped, positive  
20 stranded, RNA viruses (Lavallée et al., 1994, J. Virol.,  
~~68:1926-1934; International Patent Application No. WO~~  
90/158,75 on December 27, 1990 in the name of DANA  
FARBER CANCER INSTITUTE). These viruses code for an  
enzyme, the reverse transcriptase (RT), which enables  
25 them to replicate their RNA genome through a DNA  
intermediate. Simple retroviruses contain three,  
contiguous reading frames coding for the gag, pol and  
env genes, which constitute their structural and  
enzymatic repertoire, all packaged in the progeny  
30 virion. The gag and env genes encode the core  
nucleocapsids proteins and the membrane glycoproteins  
of the virus, respectively, whereas the pol gene gives  
rise to the reverse transcriptase and other enzymatic  
activities (ribonuclease H, integrase and protease)  
35 that are essential for viral replication. HIV belongs

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 3 -

Alamos National Laboratory, New Mexico, USA), although the open reading frame is often truncated in viral strains extensively passaged in tissue culture. The vpr open reading frame is also present in HIV-2 isolates and in most but not in all SIV isolates. A sequence similar to HIV-1 vpr is also found in Visna virus. The Vpr protein is made from a singly spliced rev-dependent mRNA species that accumulates late in infection. The Vpr protein of HIV and SIV have recently been shown to be present in mature viral particles in multiple copies. Amino acid comparison between vpx (a gene unique to HIV-2 and SIVs), and vpr from both HIV-2 and HIV-1, showed regions of strong homology (Tristem et al., 1992, EMBO J. 11: 3405-12). Like Vpr, Vpx is also packaged into the mature virion and has been shown to confer a growth advantage to viruses expressing the protein (Yu et al., 1993, J. Virol. 67: 4386-90). Interestingly, Vpx can be incorporated into HIV-1, HIV-2 or SIV with similar efficiencies. Based on the shared function, properties, including viral compartmentalization, and homologies of sequences between Vpr and Vpx, vpx in the HIV-2/SIV group, is thought to have arisen by duplication of the vpr gene (Fig. 2; Tristem et al., 1992, EMBO J. 11: 3405-12; and Myers et al., 1993, Human Retroviruses and AIDS 1993 I-II, Los Alamos National Laboratory, New Mexico, USA). Interestingly, Vpr and Vpx are the first regulatory protein of any retrovirus found to be associated with viral particles. Other regulatory proteins, such as tat, Rev, Nef, Vif and Vpu are not virion-associated. The assembly and maturation of HIV-1 viral particles is a complex process in which the structural Gag, Pol and Env gene products are expressed in the form of polyprotein precursors. The Gag proteins of HIV play

- 2 -

to the lentivirus subfamily, members of which are characterized by several additional open reading frames (ORF) not found in simple retroviruses (Fig. 1). These ORFs all appear following gag-pol sequences, either immediately preceding the env sequences or overlapping it, and at least in one case, nef, extending well into the 3' Long Terminal Repeat (LTR). These ORFs code for non-structural viral proteins readily detectable in the cells. Much evidence has accumulated indicating that these gene products, collectively referred to as auxiliary proteins, are capable of modulating viral replication and infectivity.

HIV-1 possesses at least six such auxiliary proteins, namely, Vif, Vpr, Tat, Rev, Vpu and Nef. The closely related HIV-2 does not code for Vpu, but codes for another unrelated protein, Vpx, not found in HIV-1. Mutations affecting either Tat or Rev severely impair viral replication indicating that these two auxiliary proteins are essential for viral replication. However, at least in vitro, mutations affecting other auxiliary proteins result in minimal effect on the viral replication kinetics. Hence, these proteins have been dubbed dispensable or non-essential for in vitro replication, and are usually referred to as accessory gene products.

In the past few years, it has become evident that while these "accessory" genes are not required for productive replication, they are nonetheless capable of affecting replication events, even in vitro. More importantly, recent data indicates that they may affect pathogenesis in vivo.

The vpr gene encodes a 14kDa protein (96 amino acids) in most strains of HIV-1 (Fig. 2; Myers et al., 1993, Human Retroviruses and AIDS 1993 I-II, Los

- 5 -

Interestingly, while mutations affecting HIV-1 vpr do not affect replication in peripheral blood mononuclear cells (PBMC), mutations in HIV-2 vpr results in a measurable impairment in these cells. Similarly, a recent study using anti-sense RNA directed against vpr inhibited viral replication in primary macrophages but not in transformed T-cells. Previous work indicated that this rapid growth advantage may be conferred by the weak transactivation property of Vpr on HIV-LTR directed gene expression (European Patent Application published under No. 474,797 on March 18, 1992 in the name of DANA FARBER CANCER INSTITUTE). Cotransfection experiments suggest that vpr could augment the expression of a reporter gene from several heterologous promoters by approximately three to ten fold.

The carboxyl terminal sequence of Vpr have been shown to be important for Vpr mediated transactivation as prematurely truncated proteins are non-functional and are not packaged into the virion. Interestingly, a recent report also indicated that the carboxyl terminal of the protein is important for nuclear localization (Lu et al., 1993, J. Virol., 67(1):6542-6550). A specific vpr responsive LTR sequence was not identified and the exact mechanism by which vpr augments reporter gene expression is not clear. The precise mode of action of vpr is yet to be established. However, the presence of Vpr in the viral particle (a property also shared by Vpx) suggests that this protein has a role in the early stage of infection. Virion-associated non-structural proteins in many viral systems play a pivotal enzymatic functions in early replication steps, either because cellular homologs are unavailable or are sequestered, for example, in the nucleus. It is

- 4 -

a central role in virion assembly and budding. Gag proteins are initially synthesized as myristylated polyprotein precursors, Pr55<sup>gag</sup> and Pr160<sup>gag-pol</sup>, which are transported to the inner face of the plasma membrane where they can direct particle formation, even in the absence of other viral proteins. Complete budding leads to formation of immature particles, followed by HIV protease mediated cleavage of the Gag and Gag-Pol precursor polyproteins and formation of mature HIV particles with condensed core. The mature virion proteins derived from cleavage of the gag-encoded precursor, Pr55<sup>gag</sup>, include the p17 matrix protein (MA), the p24 capsid protein (CA), the p7 nucleocapsid protein (NC), and a small proline-rich peptide of approximately 6 kDa designated p6 which are linked in this order in the polyprotein precursor. Vpr is not part of the virus polyprotein precursors and its incorporation occurs by way of an interaction with a component normally found in the viral particle.

It was recently reported that the HIV-1 Vpr could be incorporated *in trans* into viral-like particle (VLP) originating from expression of the Pr55<sup>gag</sup> only (Lavallée et al., 1994, J. Virol., 68:1926-1934). Data from this and other studies indicate that Vpr incorporation appeared to result from a direct interaction of Vpr with the carboxy-terminal region of the Pr55<sup>gag</sup> polyprotein (Paxton et al., 1993, J. Virol., 67(12):7229-7237; Lu et al., 1993, J. Virol., 67(1):6542-6550).

Functional studies indicated that the full length vpr protein could confer favorable growth properties to viruses. The increase in virion production is more pronounced in primary macrophages in both HIV-1 and HIV-2 systems, suggesting that Vpr function may be important in specific target cells.



- 7 -

HIV-2 virions to affect their structural organization and/or functional integrity.

It would also be highly desirable to be provided with a Vpr protein, a Vpx protein or  
5 fragments thereof which permit the development of chimeric molecules that can be specifically targeted into the mature HIV-1 and/or HIV-2 virions to affect their structural organization and/or functional integrity, thereby resulting in treatment of HIV-1  
10 and/or HIV-2 infections.

It would also be highly desirable to be provided with a therapeutic agent which permit the targeting of chimeric molecules into the mature HIV-1 and HIV-2 virions as a treatment for HIV-1 and HIV-2  
15 infections.

It would also be highly desirable to be provided with the identification of the protein interactions responsible for Vpr or Vpx incorporation into the mature HIV-1 and/or HIV-2 virions.

It would also be highly desirable to be provided with means to incorporate Vpr or Vpx into the mature HIV-1 and/or HIV-2 virions by making use of the protein interactions responsible for incorporation of Vpr or Vpx therein, thereby affecting the functional  
20 integrity of the HIV virions.

It would also be highly desirable to be provided with a Vpr protein fragment, a Vpx protein fragment, a p6 protein or p6 protein fragment which permits the development of molecules that can  
25 specifically interfere with the protein interactions responsible for Vpr or Vpx incorporation into the mature HIV-1 and/or HIV-2 virions to affect their functional integrity, thereby resulting in treatment of HIV-1 and/or HIV-2 infections.

- 6 -

possible that Vpr is one such protein, capable of modulating early viral specific functions such as reverse transcription stabilization of early RNA or DNA intermediates, transport to the nucleus or integration. It is equally possible that Vpr could function at an early step, in a non-viral specific manner, by triggering processes that could make the cellular environment congenial to establish viral infection. In this regard, HIV-1 Vpr has been reported to be involved in inducing cellular differentiation in rhabdomyosarcoma cells (Levy et al., 1993, Cell, 72:541-550). Finally, because Vpr is synthesized late in the infection cycle of HIV, it may regulate the morphogenesis of the virus (late events) by an unknown mechanism or constitute a structural protein involved in the integrity of the virions.

The use of transport polypeptides for biological targeting is well known and was adapted to many fields. The HIV Tat protein has been described to effect the delivery of molecules into the cytoplasm and nuclei of cells (International Application published on March 3, 1994 as No. WO 94/04686 in the name of BIOGEN, INC.). However, the Tat transport polypeptides can not allow the delivery of molecules to HIV virions. Viral proteins such as Gag of Rous sarcoma virus and Moloney murine leukemia virus and portion of HIV-1 Gag protein have been used as carrier for incorporation of foreign antigens and enzymatic markers into retroviral particles (Wang et al., 1994, Virology, 200:524-534). However, most of the Gag protein sequences are essential for efficient viral particles assembly, thus limiting the use of such virion components as carrier.

It would be highly desirable to be provided with means to target molecules to mature HIV-1 and/or

- 9 -

Another aim of the present invention is to provide a Vpr protein fragment, a Vpx protein fragment, a p6 protein or p6 protein fragment which permit the development of molecules that can specifically interfere with the protein interactions responsible for Vpr or Vpx incorporation into the mature HIV-1 and/or HIV-2 virions to affect their functional integrity, thereby resulting in treatment of HIV-1 and/or HIV-2 infections.

Another aim of the present invention is to provide a therapeutic agent which interferes with the protein interactions responsible for Vpr or Vpx incorporation into the mature HIV-1 and/or HIV-2 virions as a treatment for HIV-1 and/or HIV-2 infections.

In accordance with the present invention there is provided a protein for targeting into a mature HIV-1 and/or HIV-2 virion, the protein comprising a sufficient number of amino acids of a Vpr protein, a Vpx protein, functional derivatives or fragments thereof, wherein the protein interacts with a Gag-precursor protein of the mature virion and is incorporated by the virion. More specifically, the protein interacts with the protein p6 which is a component of the Gag-precursor protein.

More specifically, the protein of the present invention, further comprises a protein fragment covalently attached to its N- or C-terminal to form a chimeric protein which is also incorporated by the mature virion. Such an attached protein fragment of the present invention consists of amino acid sequence effective in reducing HIV expression or replication, the amino acid sequence encoding for example a RNase activity, protease activity, creating steric hindrance during virion assembly and morphogenesis and/or

- 8 -

It would also be highly desirable to be provided with a therapeutic agent which interferes with the protein interaction responsible for Vpr or Vpx incorporation in the mature HIV-1 and/or HIV-2 virions as a treatment for HIV-1 and/or HIV-2 infections.

#### DISCLOSURE OF THE INVENTION

One aim of the present invention is to provide means to target molecules to mature HIV-1 and/or HIV-2 virions to affect their structural organization and/or functional integrity

Another aim of the present invention is to provide a Vpr protein, a Vpx protein or fragments thereof which permit the development of chimeric molecules that can be specifically targeted into the mature HIV-1 and/or HIV-2 virions to affect their structural organization and/or functional integrity, thereby resulting in treatment of HIV-1 and/or HIV-2 infections.

Another aim of the present invention is to provide a therapeutic agent which permits the targeting of chimeric molecules into the mature HIV-1 and/or HIV-2 virions as a treatment for HIV-1 and/or HIV-2 infections.

Another aim of the present invention is to provide the identification of the protein interactions responsible for Vpr or Vpx incorporation into the mature HIV-1 and/or HIV-2 virions.

Another aim of the present invention is to provide means to incorporate Vpr or Vpx into the mature HIV-1 and/or HIV-2 virions by making use of the protein interactions responsible for incorporation of Vpr or Vpx therein, thereby affecting the functional integrity of the HIV virions.

- 11 -

carrier. The administration step of the method is effected intracellularly for anti-viral treatment including gene therapy or intracellular immunization of the patient through DNA transfection or  
5 administration of the chimeric protein. The anti-viral treatment can be effected through transfection of a patient's hematopoietic cells with a DNA construct harboring a Vpr/Vpx chimeric protein, followed by readministration of the transfected cells,  
10 and/or through administration of a DNA construct harboring a Vpr/Vpx chimeric protein or directly by administration of a Vpr/Vpx chimeric protein, via the blood stream or otherwise.

In accordance with the present invention there  
15 is also provided A vector comprising: (a) a DNA segment encoding a protein which interferes with the incorporation of native Vpr/Vpx into HIV-1 and/or HIV-2 virions, comprising a sufficient number of amino acids of a Vpr protein, a Vpx protein, a p6 protein,  
20 functional derivatives or fragments thereof; and (b) a promoter upstream of the DNA segment.

Accordingly, in view of two different approaches of the present invention, therapeutic agents which may be used in accordance with the  
25 present invention are selected from the group consisting of a protein of the present invention including Vpr/Vpx chimeras comprising RNase, proteases or amino acids sequences capable of creating steric hindrance during virion morphogenesis, and/or  
30 affecting viral protein interactions responsible for infectivity and/or viral replication, Vpr/Vpx protein fragment, p6 protein and p6 protein fragment, and DNA sequences encoding a protein of the present invention.

In accordance with the present invention there  
35 is also provided a pharmaceutical composition for

- 10 -

affecting viral protein interactions responsible for infectivity and/or viral replication.

More specifically, the protein of the present invention, further comprises a molecule to form a protein-molecule complex which is also incorporated by the mature virion. Such a molecule is selected from the group consisting of anti-viral agents, RNases, proteases, and amino acid sequences capable of creating steric hindrance during virion assembly and morphogenesis. The molecule of the protein-molecule complex of the present invention affects the structural organization or functional integrity of the mature virion by steric hindrance or enzymatic disturbance of the virion.

In accordance with the present invention there is also provided a protein which interferes with Vpr or Vpx incorporation into HIV-1 and/or HIV-2 virions and which comprises a sufficient number of amino acids of a Vpr protein fragment, a Vpx protein fragment, a p6 protein, p6 protein fragment, or its functional derivative thereof, wherein the protein interacts either with a Gag-precursor or with Vpr or Vpx protein to compete with the Vpr-Gag-precursor or Vpx-Gag-precursor interaction and consequently to interfere with the incorporation of the native Vpr or Vpx into the virions and to substantially prevent replication of the mature virion.

In accordance with the present invention there is also provided a method of substantially reducing HIV expression or replication in a patient infected with HIV-1 and/or HIV-2, which comprises administering at least one therapeutic agent selected from the group consisting of the protein or DNA sequences encoding the protein of the present invention, to the patient in association with a pharmaceutically acceptable

Fig. 10 shows the effect of VprIE (as shown in Fig. 9) expression on the growth rate and viability of Jurkat cells when challenged with HIV (pHxBRU or pHxBRU-R ATG<sup>-</sup>).;

5 Fig. 11 shows the effects of VprIE expression on HIV-1 replication, in stably transfected Jurkat cells; and

Fig. 12 shows the infectivity potential of HIV virions harboring a Vpr-chimeric protein; and

10 Fig. 13 shows an expression construct for the specific expression of a Vpr-based chimeric protein in HIV-infected cells.

#### MODES FOR CARRYING OUT THE INVENTION

15 In accordance with the present invention, two different approaches using the Vpr/Vpx protein and p6 protein are described herein for the treatment of HIV-1 and HIV-2 infections.

20 In the first approach, the region of Vpr/Vpx protein, which is involved in the protein interaction responsible for Vpr/Vpx virion-incorporation, is used as a carrier to target molecules to mature HIV-1 and HIV-2 virions.

25 In the second approach, the region of Vpr/Vpx protein or alternatively the region of Gag-precursor, which are both involved in the protein interaction responsible for Vpr/Vpx virion-incorporation, is used to interfere with the native viral Vpr/Vpx protein incorporation.

30 In general, the abbreviations used herein for designating the amino acids are based on recommendations of the IUPAC-IUB Commission on Biochemical Nomenclature (Biochemistry, 1972, 11:1726-1732).

- 12 -

reducing HIV expression in a patient infected with HIV-1 and/or HIV-2, which comprises a sufficient amount of the therapeutic agent of the present invention in association with a pharmaceutically acceptable carrier.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the HIV genetic organization, where the vpr gene is positioned in the central region;

Fig. 2 shows the amino acid sequences of native Vpr protein from HIVLAI isolate and native Vpr and Vpx proteins from HIV2ROD isolate;

Fig. 3 shows the mutagenesis chart of the vpr gene of HxBH10 or pHxBRU template of HIV-1;

Figs. 4A to 4E show the analysis of Vpr present in lysates and supernatants of MT4 cells infected with HIV-1 containing various mutated or truncated vpr mutants;

Figs. 5A and 5B show the stability of the truncated or mutated Vpr protein in infected MT4 cells;

Fig. 6 shows quantification of the incorporation of different mutated Vpr into HIV-1 virions;

Figs. 7A and 7B show the expression plasmids and the P6 construct plasmid expressing P6 mutant;

Figs. 8A and 8B show the incorporation of Vpr into virions in the presence of different truncated or mutated Gag P6 protein;

Fig. 9 shows the construction of a retroviral vector (pBaBepuro) for the expression of a chimeric protein between Vpr (from HIV-1 LAI strain) and a portion of a predetermined protein (pBaBepuro-VprX), such as HIV-1 Vpu protein (pBaBepuro-VprIE) or CAT (pBaBepuro-VprCAT);



- 15 -

His Phe Pro Arg Ile Trp Leu His Gly Leu Gly Gln His Ile Tyr Glu  
                   35                                  40                                  45  
 5 Thr Tyr Gly Asp Thr Trp Ala Gly Val Glu Ala Ile Ile Arg Ile Leu  
                   50                                  55                                  60  
 Gln Gln Leu Leu Phe Ile His Phe Arg Ile Gly Cys Arg His Ser Arg  
  65                                  70                                  75                                  80  
 10 Ile Gly Val Thr Gln Gln Arg Arg Ala Arg Asn Gly Ala Ser Arg Ser  
                                   85                                  90                                  95  
   (SEQ ID NO:1);  
 15 Met Ala Glu Ala Pro Thr Glu Leu Pro Pro Val Asp Gly Thr Pro Leu  
  1                                  5                                  10                                  15  
 Arg Glu Pro Gly Asp Glu Trp Ile Ile Glu Ile Leu Arg Glu Ile Lys  
                   20                                  25                                  30  
 20 Glu Glu Ala Leu Lys His Phe Asp Pro Arg Leu Leu Ile Ala Leu Gly  
                   35                                  40                                  45  
 Lys Tyr Ile Tyr Thr Arg His Gly Asp Thr Leu Glu Gly Ala Arg Glu  
  50                                  55                                  60  
 25 Leu Ile Lys Val Leu Gln Arg Ala Leu Phe Thr His Phe Arg Ala Gly  
  65                                  70                                  75                                  80  
 Cys Gly His Ser Arg Ile Gly Gln Thr Arg Gly Gly Asn Pro Leu Ser  
 30                                  85                                  90                                  95  
 Ala Ile Pro Thr Pro Arg Asn Met Gln  
                   100                                  105  
   (SEQ ID NO:2); and

---

35 Met Thr Asp Pro Arg Glu Thr Val Pro Pro Gly Asn Ser Gly Glu Glu  
  1                                  5                                  10                                  15  
 Thr Ile Gly Glu Ala Phe Ala Trp Leu Asn Arg Thr Val Glu Ala Ile  
                   20                                  25                                  30  
 40 Asn Arg Glu Ala Val Asn His Leu Pro Arg Glu Leu Ile Phe Gln Val  
                   35                                  40                                  45  
 Trp Gln Arg Ser Trp Arg Tyr Trp His Asp Glu Gln Gly Met Ser Glu  
 45                  50                                  55                                  60  
 Ser Tyr Thr Lys Tyr Arg Tyr Leu Cys Ile Ile Gln Lys Ala Val Tyr  
  65                                  70                                  75                                  80  
 50 Met His Val Arg Lys Gly Cys Thr Cys Leu Gly Arg Gly His Gly Pro  
                                   85                                  90                                  95  
 Gly Gly Trp Arg Pro Gly Pro Pro Pro Pro Pro Pro Gly Leu Val  
                   100                                  105                                  110  
 55  (SEQ ID NO:3);

functional fragments or derivatives thereof, wherein  
the Vpr/Vpx protein, its fragments or derivatives

- 14 -

From the specification and appended claims, the term therapeutic agent should be taken in a broad sense so as to also include a combination of at least two such therapeutic agents. Further, the DNA segments or proteins according to the present invention can be introduced into individuals in a number of ways. For example, erythropoietic cells can be isolated from the afflicted individual, transformed with a DNA construct according to the invention and reintroduced to the afflicted individual in a number of ways, including intravenous injection. Alternatively, the DNA construct can be administered directly to the afflicted individual, for example, by injection in the bone marrow. The DNA construct can also be delivered through a vehicle such as a liposome, which can be designed to be targeted to a specific cell type, and engineered to be administered through different routes.

In accordance with the first approach of the present invention, there is provided the use of the Vpr or Vpx protein, which is referred to as the "Vpr/Vpx" protein, functional derivatives or fragments thereof for the targeting of molecules to the HIV-1 and/or HIV-2 virions. The sequence responsible for such targeting is termed herein the Vpr/Vpx incorporation domain.

The preferred Vpr/Vpx protein, which is used in accordance with the first approach of the present invention, contains a sufficient number of amino acids corresponding to at least one of the following amino acid sequences consisting of:

Met	Glu	Gln	Ala	Pro	Glu	Asp	Gln	Gly	Pro	Gln	Arg	Glu	Pro	His	Asn
1				5				10						15	
Glu	Trp	Thr	Leu	Glu	Leu	Leu	Glu	Glu	Leu	Lys	Asn	Glu	Ala	Val	Arg
20							25						30		

- 17 -

int rfer with the native Vpr/Vpx incorporation into HIV-1 and/or HIV-2 virions. Again, the Vpr/Vpx protein fragments, p6 protein, p6 protein fragments, or functional derivatives thereof have retained their ability to interact with the native Vpr/Vpx or p6 protein, respectively. The expression "functional derivatives" when used herein is intended to mean any substitutions, deletions and/or additions of amino acids that do not destroy the functionality of the Vpr/Vpx incorporation domain or the region of the p6 protein which interacts therewith.

The preferred Vpr/Vpx protein fragments which is used in accordance with the second approach of the present invention is a fragment of the following amino acid sequence consisting of:

Met	Glu	Gln	Ala	Pro	Glu	Asp	Gln	Gly	Pro	Gln	Arg	Glu	Pro	His	Asn	
1				5				10						15		
Glu	Trp	Thr	Leu	Glu	Leu	Leu	Glu	Glu	Leu	Lys	Asn	Glu	Ala	Val	Arg	
			20				25						30			
His	Phe	Pro	Arg	Ile	Trp	Leu	His	Gly	Leu	Gly	Gln	His	Ile	Tyr	Glu	
		35				40						45				
Thr	Tyr	Gly	Asp	Thr	Trp	Ala	Gly	Val	Glu	Ala	Ile	Ile	Arg	Ile	Leu	
	50					55					60					
Gln	Gln	Leu	Leu	Phe	Ile	His	Phe									
65					70											(SEQ ID NO:4)

wherein said fragment is a region of the Vpr/Vpx protein which binds to a Gag-precursor.

The preferred p6 protein which is used in accordance with the second approach of the present invention contains a sufficient number of amino acids corresponding to the following amino acid sequences consisting of:

Leu	Gln	Arg	Ser	Pro	Glu	Pro	Thr	Ala	Pro	Pro	Glu	Glu	Ser	Phe	Arg	
1				5				10						15		
Ser	Gly	Val	Glu	Thr	Thr	Thr	Pro	Pro	Gln	Lys	Gln	Glu	Pro	Ile	Asp	
		20					25						30			

- 16 -

thereof have retained the virion-incorporation function of the native Vpr/Vpx protein.

The expression "functional fragments or derivatives" when used herein is intended to mean any  
5 substitutions, deletions and/or additions of amino acids that do not affect the virion-incorporation function of the Vpr native protein.

In accordance with the first approach of the present invention, a preferred Vpr/Vpx chimeric  
10 protein comprises an amino acid sequence of a Vpr/Vpx protein or a functional derivative thereof and a molecule attached to the amino acid sequence. Said molecule may be covalently attached at the N- or C-terminal of the amino acid sequence or it may be  
15 attached to the amino acid sequence at any amino acid position by chemical cross-linking or by genetic fusion.

A preferred molecule used in accordance with the present invention may be selected from the group  
20 consisting of an anti-viral agent and/or a second amino acid sequence which contains a sufficient number of amino acids corresponding to RNases, proteases, or any protein capable of creating steric hindrance during virion morphogenesis and/or affecting viral  
25 protein interactions responsible for infectivity and/or viral replication.

The Vpr/Vpx chimeric protein in accordance with the first approach of the present invention may be used for the targeting of molecules into the mature  
30 virions of HIV-1 and/or HIV-2, such as polypeptides, proteins and anti-viral agents, among others.

In accordance with the second approach of the present invention, there is provided the use of Vpr/Vpx protein fragments, p6 protein, p6 protein  
35 fragment, or functional derivatives thereof which

- 19 -

# HIV-1 vpr regions associated with viral particles incorporation

The substitution mutations and deletions of vpr were generated by site-directed mutagenesis. Wildtype  
5 vpr sequence and location of predicted alpha-helix structures are indicated at the top of Fig. 3. Oligonucleotide-directed mutagenesis of the vpr gene was carried out on DNA fragments derived from the pHxBRU template (Fig. 3) and then cloned into an  
10 infectious provirus (pHxBRU) (Lavallée et al., 1994, J. Virol., 68:1926-1934). Amino acid substitutions are indicated. BRUR 77/79, BRUR 76/79 and HxBH10 72/78 (Yao et al., 1992, J. Virol. 66:5119-5126) are truncated vpr proteins with additional unrelated amino  
15 acids generated by frame shift mutations.

Figs. 4A-4E are autoradiograms that illustrate the analysis of truncated or substituted Vpr mutants in HIV-1 infected MT4 cells.  $2 \times 10^6$  T-lymphoid cells (MT4) were infected (or transfected, Figs. 4D & 4E)  
20 with HIV-1 which contain wildtype or truncated Vpr (Fig. 3), as shown at the top of the autoradiograms.

The position of HIV-1 viral proteins are indicated at the left of the autoradiograms (Vpr). At 40 h post-transfection, cells were labelled with 100  $\mu$ Ci of  $^{35}$ S-methionine and 100  $\mu$ Ci of  $^3$ H-leucine for 16 h.  
25 Virions were pelleted from cell-free supernatants by ultracentrifugation at 35,000 rpm through a 20% sucrose cushion for 2 h. Both cells (Figs. 4A, 4C [left] & 4D) and sucrose cushion pelleted viruses  
30 (Figs. 4B, 4C[right] & 4E) were lyzed in RIPA buffer (140 mM NaCl, 8 mM NaHPO<sub>4</sub>, 2 mM NaH<sub>2</sub>PO<sub>4</sub>, 1% Nonidet<sup>TM</sup> P-40, 0.5% sodium deoxycholate, 0.05% SDS) and immuno-precipitated with a HIV-1 positive human serum combined with a rabbit anti-Vpr serum. Proteins were  
35 then analyzed on a sodium dodecyl sulfate (SDS) 12.5%-17% gradient polyacrylamide gel electrophoresis (PAGE)

- 18 -

Lys Glu Leu Tyr Pro Leu Thr Ser Leu Arg Ser Leu Phe Gly Asn Asp  
 35 40 45

Pro Ser Ser Gln  
 5 50

(SEQ ID NO:5).

The Vpr/Vpx fragment, p6 protein and p6 protein fragment in accordance with the second approach of the present invention may be used for interfering with the virion-incorporation of native Vpr/Vpx into HIV-1 and/or HIV-2 virions.

The purpose of the treatment in accordance with the first and second approaches of the present invention may be a prevention or a treatment. The product in these treatment procedures may be expressed intracellularly or provided to the cell via the blood stream.

In accordance with the first approach of the present invention, the expressed or administered product may be effective in the production of defective viral particles, for instance, viral particles with Vpr/Vpx chimeric proteins such as the ones associated with virally directed protease or nuclease or with a portion of protein which affects the structural organization and/or functional integrity of the virions.

The treatment in accordance with the second approach of the present invention may consists in the production of viral particles having substantially reduced replication capacity, for instance, HIV-1 and HIV-2 viral particles devoided of functional level of Vpr/Vpx protein as a consequence of Vpr-Gag-precursor or Vpx-Gag-precursor interaction interference using Vpr/Vpx protein fragments, p6 protein and p6 protein fragments.

35

- 21 -

## HIV-1 GAG P6 regions associated with Vpr incorporation

To investigate the mechanism of incorporation of Vpr, the ability of Gag-expressor plasmids, harboring deletions or mutations in the C-terminus of the capsid precursor, to target Vpr into virions in transiently transfected cells was tested. In this system two expressor plasmids described in Fig. 7A were cotransfected into COS-7 cells (Lavallée et al., 1994, J. Virol., 68:1926-1934; Fig. 7A). Deletions are shown as dotted lines between the thick lines. ptrENV contains 3109 (nucleotides 989 to 4098) and 1294 (nucleotides 5925 to 7219) base pair deletions affecting respectively gag, pol and the gp120 domain of env genes. ptrENV encodes Vpr as well as all HIV-1 auxilliary proteins (Vif, Tat, Rev, Vpu Nef and gp41). The pIIIgagCAR plasmid, a rev-dependent Gag expressor, which encodes Pr55<sup>gag</sup> and the protease domain of the pol gene (PR), contains the Rev-responsive element (RRE/CAR) sequence. P6 is the C-terminal components of the Pr55<sup>gag</sup> precursor (Fig. 7A). Fig. 7B illustrates the P6 constructs. Plasmids expressing P6 mutant were generated by introducing a termination codon or a substitution by polymerase chain reaction (PCR)-based site-directed mutagenesis in pIIIgagCAR plasmid.

Figs. 8A & 8B illustrate the trans incorporation of Vpr into virus-like particles. COS-7 cells were transfected with pIIIgagCAR plasmid (lane 1), or ptrENV plasmid (lane 2) or cotransfected with both constructs (lane 3). ptrENV was cotransfected with pIIIgagCAR based construct harboring a substitution or a premature termination codon in the p6 protein: L1/stop (lane 4), S17/stop (lane 5), Y36/stop (lane 6), P10,11L (lane 7), L44/stop (lan 8), P49/stop (lane 9). 48 h posttransfection [<sup>35</sup>S]methionine- and

and autoradiography. Quantification of virion associated Vpr and protein stability was determined by densitometric scanning of the autoradiograms using a laser densitometer (Molecular Dynamics™ densitometer).

5           Fig. 5 shows the stability of different truncated (Fig. 5B) or substituted (Fig. 5A) Vpr mutants in HIV-1 infected cell lysates. The intensity of wildtype or mutated Vpr proteins were quantified relative to the intensity of the p66 reverse  
10 transcriptase (RT) bands. Immunoprecipitation analysis has shown that all truncated Vpr proteins were present at low level in cell lysates suggesting the importance of the C-terminal region for Vpr stability (Figs. 4A and 5B).

15           Fig. 6 shows the efficiency of incorporation of different mutated Vpr into HIV-1 virions. The incorporation of mutated Vpr into virions was also evaluated by densitometric analysis. The intensity of Vpr proteins into virions were quantified relative to  
20 the intensity of the p66 reverse transcriptase (RT)

---

bands in autoradiograms presented in Figs. 4C and 4E. The results of the present invention demonstrate that substitution mutations (A30F, H33L and E25K) in the N-terminal portion of Vpr significantly impair the  
25 incorporation of Vpr protein into virions (Figs. 4C, 4E & 6). It should be noted that this region of the protein is predicted to form an alpha helix and thus is reminiscent of a structure involved in protein-protein interaction. Interestingly, an alpha helix is  
30 also predicted from the corresponding region of Vpx. These data indicate that the N-terminus of Vpr is important for Vpr incorporation in the virion. This region will be further defined by analyzing additional Vpr mutants for their stability and virion-  
35 incorporation capacity.



- 23 -

Prokaryotic expression are useful for the preparation of large quantities of the protein encoded by the DNA sequence of interest. This protein can be purified according to standard protocols that take advantage of the intrinsic properties thereof, such as size and charge (i.e. SDS gel electrophoresis, gel filtration, centrifugation, ion exchange chromatography...). In addition, the protein of interest can be purified via affinity chromatography using polyclonal or monoclonal antibodies. The purified protein can be used for therapeutic applications.

For administration to humans, the prescribing medical professional will ultimately determine the appropriate dosage for a given patient, and this can be expected to vary according to the chosen therapeutic regimen (i.e DNA construct, protein, cells), the response and condition of the patient as well as the severity of the disease.

The present invention will be more readily understood by referring to the following examples which are given to illustrate the invention rather than to limit its scope.

25

EXAMPLE I

Incorporation of a specific epitope into retroviral particles from a HIV-1 cloned provirus (cis incorporation)

30

The ability of a foreign antigen to be incorporated when expressed in cis (with respect to the viral genome) into viral particles when fused to Vpr protein was examined. Trans-incorporation (trans with respect to the viral genome) of Vpr chimeric proteins into Vpr-virion in MT4 cells presents

- 22 -

[<sup>3</sup>H]leucine-labelled viral proteins were immuno-precipitated, from the cell lysates (Fig. 8A) or the cell-free supernatant (Fig. 8B) centrifuged through a 20% sucrose cushion, with the HIV-1 positive human serum 162 mixed with a rabbit anti-Vpr polyclonal antibodies and analysed by SDS-PAGE and autoradiography.

The 14kDa vpr product can be detected in the pelleted virions produced by cells cotransfected with pIIIgagCAR and ptrENV or the P49/stop or P10,11L mutants (Fig. 8B, lanes 3, 9 and 7, respectively). However, virions produced from cells cotransfected in the presence of ptrENV and L1/stop, S17/stop, Y36/stop or L44/stop constructs lacked detectable Vpr (Fig. 8B, lanes 4, 5, 6 and 8, respectively). These results indicate a direct correlation between the absence of p6 and the loss of Vpr incorporation, suggesting that p6 is directly implicated. Moreover, deletion analysis suggests that the carboxyl terminal of p6 is important for Vpr incorporation. It has already been mentioned that Vpr and Vpx share strong regions of homology. Strikingly, the region of highest homology is found in the region of Vpr which has been shown to interact with p55Gag. This region which is predicted to form an alpha helical structure is thought to be the packaging region of Vpr/Vpx.

Once known, these incorporation domains can be used in a variety of ways. For example, fusion proteins thereof can be inserted in expression vectors according to standard procedures. Such vectors contain all the necessary regulatory signals to promote expression of the fusion protein of interest. Typically, expression vectors are prokaryote specific or eukaryote specific although shuttle vectors are also widely available.

- 25 -

Vpr) antisera are able to immunoprecipitate the chimeric product from the virions. These data indicate that the Vpu epitope was successfully transferred into virions when expressed as a Vpr fusion product.

5

#### EXAMPLE II

#### Expression of Vpr chimeras in trans from retroviral vectors

10 Since one purpose of the present invention is to provide a therapeutic and/or prophylactic agent for HIV infections, an in vitro model was developed in order to express a vpr/vpx-chimeric protein in trans with respect to the viral genome. The ability of a  
15 foreign antigen to be incorporated in trans into viral particles was examined then with this in vitro model.

#### **A-Construction of retroviral vectors encoding Vpr fusion proteins**

20 Chimeric molecules were developed by fusion of Vpr sequences with either sequences encoding different enzymatic activities or random amino acid sequences of different lengths. In addition to the already described VprIE (Fig. 9), Vpr was fused to the  
25 prokaryotic gene chloramphenicol acetyl-transferase (CAT), to generate VprCAT (Fig. 9). To generate stable CD4<sup>+</sup> T cell lines expressing constitutively Vpr-based chimeric proteins, retroviral vector technology was used. Since an important feature in  
30 any gene therapy approach is the method by which foreign DNA is introduced into cells, the method used to transduce a desired gene inside the cells must be accurately chosen. Retroviral vectors are preferred vectors for gene transfer of mammalian cells in  
35 culture, because they lead to high efficiency

limitations due to the low transfection efficiency-  
obtained when two expression vectors are transfected  
in T cell lines. To overcome these limitations, a  
construct expressing a Vpr chimeric protein was  
5 designed. A Vpr-Vpu fusion protein was cloned into a  
Vpu-minus HIV-1 provirus plasmid (pHxBRU; Lavallée et  
al., 1994, J. Virol., 68:1926-1934), the fusion pro-  
tein (VprIE) contains the first 88 amino acids of Vpr  
from the LAI strain and a Vpu epitope (Fig. 9). The  
10 last 18 C-terminal amino acids of Vpu which contains a  
specific epitope (Cohen et al., 1988, Nature, 334:532-  
534) has been cloned into a XbaI-created restriction  
site located at the 3' end of the Vpr sequence, at  
position 5410 on the pHxBRU genome to yield pHxBRU-  
15 VprIE. The Vpu epitope is recognized by the specific  
Rabbit anti-Vpu peptide serum described by (Cohen et  
al., 1988, Nature, 334:532-534). The pHxBRU-VprIE  
construct was transfected in MT4 cells and the  
incorporation of the Vpr-Vpu fusion protein into  
20 virions was measured.

Two million MT4 cells were transfected with the  
pHxBRU VprIE provirus. 48 h post-transfection, cells  
were labelled with 100  $\mu$ Ci of  $^{35}$ S-methionine and 100  
 $\mu$ Ci of  $^3$ H leucine for 16 h. Virions were pelleted  
25 from the cell-free supernatants by ultracentrifugation  
at 35,000 rpm through a 20% sucrose cushion for 2 h.  
Both cells and sucrose cushion pelleted viruses were  
lyzed in RIPA buffer (140 mM NaCl, 8 mM NaHPO<sub>4</sub>, 2 mM  
NaH<sub>2</sub>PO<sub>4</sub>, 1% Nonidet<sup>™</sup> P-40, 0.5% sodium deoxycholate,  
30 0.05% SDS) and immunoprecipitated with a HIV-1 posi-  
tive human serum combined with a rabbit anti-Vpr serum  
or with a rabbit anti-Vpu serum. Analysis of vpr  
products in the cell lysates and supernatants has  
revealed that 1) Vpr-Vpu chim ras are stably expressed  
35 in transfected cells and 2) both Vpu ( $\alpha$ Vpu) and Vpr ( $\alpha$



- 27 -

VprX, cells were seeded in 100-mm petris dishes for 48 hours. Vpr-X indicates VprIE, VprCAT or any Vpr (or Vpx) chimeric protein. The medium was then removed and replaced by 4 ml of fresh medium. 16 hours later  
5 the medium containing a high level of retroviruses was harvested and used to infect  $3 \times 10^6$  of centrifuged Jurkat cells. After 48 hours, the transduced Jurkat cells were cultured in fresh medium containing 0,2 ug/ml puromycin for 3 weeks. Jurkat-VprCAT and Jurkat-  
10 VprIE resistant cell lines were then isolated and amplified.

C- Expression of Vpr-based chimeric proteins in Jurkat-VprX cell lines.

15 In the stable Jurkat cell line described in (B), the fusion protein derived from pBaBepuro-VprX are driven by the 5' MoMuLV LTR and are expected to be constitutively expressed. To evaluate the expression of the chimeric proteins and to demonstrate that large  
20 molecules and functional enzymatic activities can be efficiently transferred to the virion, the CAT activity associated with the virions generated in transduced Jurkat cells was measured. The rapid, sensitive, quantifiable and reproducible assays have  
25 been extensively described to measure the activity of CAT (European Patent Application published under No. 474,797 on March 18, 1992 in the name of DANA FARBER CANCER INSTITUTE)

Briefly,  $3 \times 10^6$  Jurkat-VprCAT cells were  
30 infected with equivalent amounts of HIV-1 Vpr+ or Vpr- virus (pHxBRU and pHxBRU-RATG<sup>-</sup>, respectively), as evaluated by reverse transcriptase (RT) activity associated with the HIV-1 Vpr+ or Vpr- viral stocks (100 000 CPM), in 1 ml of culture medium for 3 hours  
35 and then diluted in 6 ml of medium (Aldovini t al.,

- 26 -

transformation of mammalian cells through stable integration of proviral DNA into transcriptionally active regions of cellular genome. These vectors contain cis-acting elements [long terminal repeats (LTR), primer binding site (PBS), packaging signal (psi), and polyuridine track], required for viral replication, integration, gene expression and packaging as well as a selectable marker. The transacting proteins (Gag, Pol and Env), required for viral assembly, infection, replication and integration are provided by helper plasmid(s) present in packaging cell lines and are expressed from an RNA lacking a psi signal.

The complete Vpr-CAT and Vpr-IE gene were amplified by PCR with mutagenic BamHI oligonucleotides and inserted into the unique BamHI restriction site of the retroviral vector pBaBepuro, which contains a puromycin resistant gene (Fig 9; Morgenstern et al., 1990, Nucl. Acids Res. 18: 3587-96). pBaBepuro-VprCAT and pBaBepuro-VprIE were then transfected into the amphotropic Damp packaging cell line. After selection with puromycin, Damp cell lines (Damp-Vpr-CAT and Damp-Vpr-IE) able to produce recombinant retroviruses (Morgan et al., 1993, Annu. Rev. Biochem. 62: 191-217) were generated.

#### B-Generation of Jurkat CD4<sup>+</sup> T cell lines expressing constitutively the VprCAT and VprIE proteins

The amphotropic murine retroviruses encoding pBaBepuro-VprCAT or pBaBepuro-VprIE (Fig. 9), produced respectively from Damp-VprCAT and Damp-VprIE, were used to transduce CD4<sup>+</sup> Jurkat cells (CD4<sup>+</sup> T cell lines highly permissive to HIV infection). Following puromycin selection Jurkat-VprCAT and Jurkat-VprIE cell lines were isolated. Briefly,  $1,2 \times 10^6$  Damp-

- 29 -

Vpr+ strain (pHxBRU) suggests a competition for virion incorporation between the wild type Vpr and the VprCAT chimeric protein.

These data indicate that the Vpr fusion protein  
5 can be successfully expressed in transduced cells and transferred into virions when expressed in trans from a CD4<sup>+</sup> T cell line. In addition the data demonstrate that enzymatic activity associated with the chimeric protein can be targeted into HIV virions. Thus, as  
10 shown previously for the Vpr/Vpx chimera expressed from a recombinant HIV provirus, the VprCAT chimera can be incorporated in trans in HIV virions when expressed from a stable cell line.

15

EXAMPLE III

Vpr-based chimeric proteins and protection against HIV  
infection

The in vitro model described above, which  
20 expresses constitutively Vpr chimeras, was used to test the antiviral activity of these fusion proteins on viral replication.

A- Characterization of Jurkat cell lines expressing  
25 Vpr chimeric proteins

To demonstrate that foreign molecules and functional enzymatic activities can be expressed in CD4<sup>+</sup> Jurkat cell lines without affecting cell physiology, cell growth in non-infected Jurkat-VprX  
30 cell lines was analyzed. During 5 weeks, growth was monitored by assessing cell viability using the Trypan Blue exclusion method (every three days; Aldovini et al., Eds, 1990, Techniques in HIV research, Stockton Press). All the Jurkat-VprX cell lines were shown to  
35 grow at a rate similar to the parental Jurkat cell

Eds, 1990, Techniques in HIV research, Stockton Press). pHxBRU-RATG<sup>-</sup> was generated by introducing a GTG codon instead of the ATG initiation codon in the Vpr gene from the pHxBRU proviral clone (Lavallée et al., 1994, J. Virol., 68:1926-1934). Following viral adsorption, cells were passed every three days. After 20 days, during the peak of viral production, supernatants containing the virus were collected. Virions were then purified by ultracentrifugation through a 20% sucrose cushion (Lavallée et al., 1994, J. Virol., 68:1926-1934). The pelleted virus were resuspended in 80 ul of Tris-HCl 250 mM and lysed by freeze and thaw. 70 ul of the lysed virus was then assayed for CAT activity. Table 1 shows the CAT activity associated with the HIV-1 virions produced in the Jurkat VprCAT transduced cells.

TABLE 1

CAT activity associated to HIV-1 virions produced in a Jurkat-VprCAT transduced cell line

	CAT Activity
	% Acetylation
Mock	0,29
Jurkat-VprCAT/pHxBRU	1,44
Jurkat-VprCAT/pHxBRU-R ATG <sup>-</sup>	5,09

Detection of CAT activity in the virions thus revealed that 1) the VprCAT chimera is constitutively expressed in transduced Jurkat cells, 2) the fusion protein is incorporated into viral particles following infection and 3) the lower CAT activity detected in virus produced from transduced cells infected with the HIV-1



- 31 -

infection when challenged with the HIV-1 vpr<sup>-</sup> strains. In contrast the parental Jurkat cell line exhibited full viral production 15 days post infection when challenged with either the Vpr<sup>+</sup> or Vpr<sup>-</sup> viruses. In addition, viable cell counts demonstrated that Jurkat-VprIE cells had survived HIV-1 infection even after 27 days. Indeed, as previously mentioned the rate of cellular growth was identical to the uninfected parental Jurkat cell control (figure 10). In contrast, the infected parental cell culture was completely killed by day 22. (Aldovini et al., Eds, 1990, Techniques in HIV research, Stockton Press).

The data obtained demonstrate that 1) the VprIE fusion protein when provided in trans in Jurkat cells can efficiently delay HIV-1 replication and cytopathic effect such as syncytium formation; and 2) the VprIE chimeric protein was successfully able to protect Jurkat cells from HIV-1 infection for more than 25 days post-infection while allowing normal cell growth.

20

---

#### B- Incorporation of Vpr-chimeric protein into HIV virion affects viral infectivity

The assembly and maturation of HIV-1 viral particles is a complex process which involves multiple interactions between viral and cellular constituents. The presence of non-relevant amino acid sequence fused to Vpr during assembly and budding may, due to steric hindrance, affect viral morphogenesis and maturation, thereby resulting in the production of virus with drastically reduced infectious potential. Indeed, as shown above the introduction of 18 amino acids derived from HIV-1 Vpu at the C-terminal end of Vpr reduced considerably the replication and spread of HIV in tissue culture. Similar results were also obtained when a larger amino-acid sequence encoding CAT was

35

- 30 -

line (mock-Jurkat TA), as evidenced in Fig 10, by the virtual super-imposition of the growth rate curves of the parental Jurkat cell line, mock infected, and that of Jurkat-Vpr-IE/pHxBRU-R ATG<sup>-</sup>.

5       The ability of Vpr fusion proteins to protect CD4<sup>+</sup> T cell lines against HIV infection was then evaluated. The Jurkat-VprIE and Jurkat-VprCAT cell lines were challenged with HIV vpr<sup>+</sup> and HIV vpr<sup>-</sup> viruses (pHxBRU and pHxBRU-R ATG<sup>-</sup>, respectively) to  
10       assess their susceptibility to viral infection. The infection kinetic was monitored by measuring viral production (RT activity in the supernatant) and the presence of cytopathic effects (syncytium formation and viable cell counts) every three days during four  
15       weeks (Aldovini et al., Eds, 1990, Techniques in HIV research, Stockton Press). Briefly, 3 X 10<sup>6</sup> cells of each Jurkat cell lines were infected with a 100 000 CPM RT activity-associated viral stock in 1 ml of medium for 3 hrs and then diluted in 6 ml with culture  
20       medium. Every 3 days during one month, RT activities were measured from supernatant aliquots and the number of cells in the cultures was determined by counting, from aliquot therefrom, the number of cells able to exclude trypan blue dye. The cell cultures were  
25       diluted to approximately 500,000 cells per ml and 200 ul of supernatant were harvested for R.T. activity detection (Aldovini et al., Eds, 1990, Techniques in HIV research, Stockton Press). The data indicate that all CD4<sup>+</sup> T cell lines expressing Vpr chimeric proteins  
30       exhibited, albeit to a different extent, a non permissiveness to HIV infection as compared to the highly HIV permissive parental cells (mock-Jurkat TA). Figure 11 shows that the Jurkat-VprIE exhibited a peak viral production 21 days post-infection when  
35       challenged with the HIV-1 vpr<sup>+</sup> virus and 27 days post

- 33 -

fold decrease in viral infectivity can be measured in viruses that incorporated Vpr-chimeric proteins as compared to wild-type virus .

5

**EXAMPLE IV****Use of Vpr-protease chimeric proteins against HIV infections**

The examples described above demonstrated that  
10 Vpr chimeric proteins are targeted into HIV virions and affect the structural organization and/or functional integrity of the virions.

The following example will illustrate the use of  
a Vpr-antiviral chimera, such as a Vpr-protease  
15 chimera for protecting the cells, in which it is expressed, against HIV infection. Indeed, degradation of the protein content in the virions will lead to completely defective viral particles even though small level of Vpr-protease chimeric molecules are  
20 incorporated.

---

**A-Construction of Vpr-based chimeric molecules in which the Vpr incorporation domain is fused to peptide sequences exhibiting potential antiviral properties**

25 Virus-encoded RNA and proteins represent the major components of viral particles. A preferred strategy according to the present invention takes advantage of the experiments that had indicated that functional enzymatic activity can be targeted to the  
30 virion as Vpr-based chimeric molecules. Using a similar approach, a sequence encoding a proteolytic activity can be fused to the Vpr C-terminus. It is believed that the presence of such an activity in the virion will lead to the destruction and/or formation  
35 of defective viral particles. Proteases are enzymes

- 32 -

fused to Vpr and similar results are expected using a Vpx-Vpu fusion or Vpx-X fusion.

To evaluate the infectivity potential of HIV virions that had incorporated the Vpr-CAT or Vpr-IE fusion proteins, Jurkat-VprCAT and Jurkat-VprIE cells were transfected with HIV-1 vpr<sup>+</sup> or HIV-1 vpr<sup>-</sup> viruses. Briefly, 10 x 10<sup>6</sup> Jurkat or Jurkat-VprX cells were transfected by the DEAE-Dextran method with 10ug of HIV-1 vpr<sup>+</sup> (pHxBRU) or HIV-1 vpr<sup>-</sup> (pHxBRU-R ATG<sup>-</sup>) infectious molecular clones. After 12 and 24 hr, p24 assay (ELISA) was performed on culture supernatants to evaluate viral production. The level of p24 antigen was similar in all samples indicating that no impairment at the level of gene expression, assembly and release of the viral particles had occurred. These results indicated that the infectivity potential of the released particles that incorporated Vpr-based chimeric molecules may be affected. To evaluate the infectious potential of HIV virions containing Vpr chimeric proteins, a MAGGI assay was performed (Kimpton et al., 1992, J. Virol. 66: 2232-39). Briefly, 35,000 HeLa CD4<sup>+</sup> B-gal cells were seeded in 24 plates and infected 24 hr later with 300,000 CPM (by RT) of HIV-1, produced from Jurkat or Jurkat-VprX cells infected with HIV-1 vpr<sup>+</sup> or vpr<sup>-</sup> strains, respectively. 24 hours post-infection B-galactosidase (B-gal) activity expressed in infected cells was measured. The level of B-gal activity following infection represents a measure of the infectivity of the virus. Figure 12 shows that HIV-1 containing Vpr chimeric proteins are strongly impaired in their ability to infect CD4<sup>+</sup> cells. Overall the data indicates that 1) gene expression, assembly and release are not affected by the expression of Vpr-chimeric proteins in infected cells; and 2) a 5 to 10-



- 35 -

formation of inactive enzymatic products during intracellular biosynthesis. However, as one goal is the destruction or impairment of infectious viral particles, the constructs preferably contains a regulatory mechanism that can specifically activate the enzymatic activity of the protease in the virions. The HIV-1 Gag precursor proteolytic processing sites are cleaved specifically and sequentially by the viral protease during virion budding and maturation. When full-length Gag proteins are digested with recombinant HIV-1 protease in vitro, four of the five major processing sites in Gag are cleaved at rates that differ by as much as 400-fold. Preferably, it will introduce an HIV-1 specific protease cleavage site sequence, corresponding by exemple to the p17/p24 processing site (Myers et al., 1993, Human Retroviruses and AIDS 1993 I-II, Los Alamos National Laboratory, New Mexico, USA)), between the Vpr virion incorporation domain and the protease sequences (Figure 13). Based on the hydrophobic/hydrophilic property and enzymatic cleavage kinetic ( $V_{max}$ ,  $K_i$ ) of the p17/p24 processing site (intermediate rate of cleavage), this sequence is predicted: i) to be located at the surface of the fusion proteins; and ii) to be digested only within the virions since only there substrate will reach concentrations necessary for efficient proteolysis. During budding and maturation, cleavage of Vpr-protease chimeric protein by the HIV-1 protease will activate the prokaryotic enzymatic activity in the virions. It should be understood that other types of linkers between the Vpr/Vpx sequence and the chosen antiviral sequence are well known and available to the skill d artisan. The presence of such non-specific protease activity in the

that catalyze the cleavage of peptide bonds of protein substrates. Many eukaryotic and prokaryotic proteases have been extensively studied and are commercially available as reagents for molecular biology. The  
5 selection of appropriate proteases for Vpr-based gene therapy will be preferentially based on the following characteristics: i) the candidate protease has a large spectrum endopeptidase activity; ii) the protease  
10 should be functional in the physiological conditions found in the virion; and iii) its molecular weight should not interfere with virion packaging.

Uncontrolled proteolytic activity can destroy the protein component of cells and tissues. One  
15 naturally-occurring mechanism for controlling protease activity during biosynthesis is the presence of an amino-terminal extension fused to the enzyme. Proteolytic cleavage of the peptide bond between the so-called propeptide and the enzyme essentially  
20 reverses this evolutionary addition and restores enzymatic activity. It is thus a preferred embodiment of the present invention to use the Vpr/Vpx virion incorporation domain as a propeptide and fuse it to the active proteolytic domain of protease. One  
25 candidate protease that is considered for fusion to the Vpr/Vpx sequence is a prokaryotic protease such as the calcium-dependent protease of the cyanobacterium *Anabaena*, or a eukaryotic protease such as the alkaline exoprotease (ALP) of *Aspergillus fumigatus* and Cathepsin G. The advantage of prokaryotic  
30 proteases over eukaryotic proteases is that they do not require to go through an intracellular pathway for maturation and proper folding.

The presence of an amino terminal Vpr sequence (or Vpx sequence) in Vpr-protease fusion products (or  
35 its Vpx counterpart) can be designed to lead to the

- 37 -

to deliver and stably integrate therapeutic genes into human cells (Morgan et al., 1993, Annu. Rev. Biochem. 62: 191-217). In a preferred embodiment of the present invention, the gene transfer experiments are carried out by cloning the Tat-inducible minigene encoding the Vpr-based chimeric protein (Figure 13) into an appropriate Mo-MuLV-derived retroviral vector. A number of these retroviral vectors are readily available and well known in the field of molecular biology and gene therapy. As a control, a retroviral vector containing a frameshift mutation in the Vpr virion incorporation domain can also be generated. According to the well known methods of gene therapy, to produce amphotropic pseudotyped vector particles, the retroviral vector is transfected into the psi CRIP amphotropic packaging cell line to generate retroviral vector producer cell lines. Amphotropic pseudotyped retroviral vector particles capable of infecting human cells, can then be tested for: i) their titer using NIH-3T3 and human CD4<sup>+</sup>lymphocyte-derived CEM-10 (adherent) cell lines; ii) lack of recombinant virus via southern blot analysis; and iii) lack of helper virus production via infection of a mouse cell line. The Vpr-protease chimera will be tested in CD4<sup>+</sup> T cell lines or CD4<sup>+</sup> peripheral blood lymphocytes (PBL) to evaluate the ability of the fusion protein to protect the cells against HIV infection.

While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the

- 36 -

virion is predicted to lead to the destruction and/or generation of defective viral particles.

5 B-Structure of the minigene encoding Vpr-based chimeric proteins under HIV-1 Tat regulation

During gene therapy, Vpr/Vpx-chimeric proteins can be expressed in trans from a foreign gene integrated in the cell genome. In addition, expression of the Vpr-chimeric proteins will be  
10 preferentially restricted to HIV-1 infected cells since: i) long term expression of Vpr/Vpx-chimeras could lead to cellular toxicity; and ii) existing cell-mediated immunity (in patients) could lead to the death of all transformed cells expressing the anti-  
15 HIV-1 protein, although not infected by HIV-1.

Selected fusion proteins are preferentially cloned 3' to a constitutive or inducible promoter. The cytomegalovirus (CMV) early gene promoter which is an example of a powerful constitutive promoter.  
20 Numerous types of promoters could also be used. To ensure a specific control of the transgene expression, the CMV promoter is preferentially fused to the transactivator (Tat) responsive element (TAR), which in response to HIV-1 Tat binding, mediates an increase  
25 of transcription initiation and facilitates mRNA elongation (Fig. 13). Thus full expression of the transgene can be restricted to cells expressing Tat, such as in HIV-1 infected cells.

30 C-Construction of retroviral vectors that express a Vpr-based chimera

An important feature in any gene therapy approach is the method by which foreign DNA is introduced into cells. Safe replication-defective retroviral vectors  
35 have already been used in experimental clinical trials



- 39 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Universite de Montreal
- (B) STREET: 2900 Edouard- Montpetit
- (C) CITY: Montreal
- (D) STATE: Quebec
- (E) COUNTRY: Canada
- (F) POSTAL CODE (ZIP): H3T 1J4
- (G) TELEPHONE: 514 343 2307
- (H) TELEFAX: 514 343 2326

(ii) TITLE OF INVENTION: PROTEIN TARGETING INTO HIV VIRIONS BASED  
ON  
HIV-1 VPR FUSION MOLECULES

(iii) NUMBER OF SEQUENCES: 5

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/301,915
- (B) FILING DATE: 07-SEP-1994

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 96 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Met	Glu	Gln	Ala	Pro	Glu	Asp	Gln	Gly	Pro	Gln	Arg	Glu	Pro	His	Asn
1				5				10						15	
Glu	Trp	Thr	Leu	Glu	Leu	Leu	Glu	Glu	Leu	Lys	Asn	Glu	Ala	Val	Arg
			20				25						30		
His	Phe	Pro	Arg	Ile	Trp	Leu	His	Gly	Leu	Gly	Gln	His	Ile	Tyr	Glu
			35				40						45		
Thr	Tyr	Gly	Asp	Thr	Trp	Ala	Gly	Val	Glu	Ala	Ile	Ile	Arg	Ile	Leu
			50			55					60				
Gln	Gln	Leu	Leu	Phe	Ile	His	Phe	Arg	Ile	Gly	Cys	Arg	His	Ser	Arg
65				70						75				80	
Ile	Gly	Val	Thr	Gln	Gln	Arg	Arg	Ala	Arg	Asn	Gly	Ala	Ser	Arg	Ser
				85					90					95	

- 38 -

art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as follows in the scope of the appended claims.

- 41 -

Ser Tyr Thr Lys Tyr Arg Tyr Leu Cys Ile Ile Gln Lys Ala Val Tyr  
 65 70 75 80  
 Met His Val Arg Lys Gly Cys Thr Cys Leu Gly Arg Gly His Gly Pro  
 85 90 95  
 Gly Gly Trp Arg Pro Gly Pro Pro Pro Pro Pro Pro Gly Leu Val  
 100 105 110

## (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 72 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Glu Gln Ala Pro Glu Asp Gln Gly Pro Gln Arg Glu Pro His Asn  
 1 5 10 15  
 Glu Trp Thr Leu Glu Leu Leu Glu Glu Leu Lys Asn Glu Ala Val Arg  
 20 25 30  
 His Phe Pro Arg Ile Trp Leu His Gly Leu Gly Gln His Ile Tyr Glu  
 35 40 45  
 Thr Tyr Gly Asp Thr Trp Ala Gly Val Glu Ala Ile Ile Arg Ile Leu  
 50 55 60  
 Gln Gln Leu Leu Phe Ile His Phe  
 65 70

## (2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 52 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Leu Gln Arg Ser Pro Glu Pro Thr Ala Pro Pro Glu Glu Ser Phe Arg  
 1 5 10 15  
 Ser Gly Val Glu Thr Thr Thr Pro Pro Gln Lys Gln Glu Pro Ile Asp  
 20 25 30  
 Lys Glu Leu Tyr Pro Leu Thr Ser Leu Arg Ser Leu Phe Gly Asn Asp  
 35 40 45  
 Pro Ser Ser Gln  
 50

- 40 -

## (2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 105 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

Met Ala Glu Ala Pro Thr Glu Leu Pro Pro Val Asp Gly Thr Pro Leu
 1           5           10           15
Arg Glu Pro Gly Asp Glu Trp Ile Ile Glu Ile Leu Arg Glu Ile Lys
          20           25           30
Glu Glu Ala Leu Lys His Phe Asp Pro Arg Leu Leu Ile Ala Leu Gly
          35           40           45
Lys Tyr Ile Tyr Thr Arg His Gly Asp Thr Leu Glu Gly Ala Arg Glu
          50           55           60
Leu Ile Lys Val Leu Gln Arg Ala Leu Phe Thr His Phe Arg Ala Gly
          65           70           75           80
Cys Gly His Ser Arg Ile Gly Gln Thr Arg Gly Gly Asn Pro Leu Ser
          85           90           95
Ala Ile Pro Thr Pro Arg Asn Met Gln
          100           105

```

## (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 112 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

```

Met Thr Asp Pro Arg Glu Thr Val Pro Pro Gly Asn Ser Gly Glu Glu
 1           5           10           15
Thr Ile Gly Glu Ala Phe Ala Trp Leu Asn Arg Thr Val Glu Ala Ile
          20           25           30
Asn Arg Glu Ala Val Asn His Leu Pro Arg Glu Leu Il Phe Gln Val
          35           40           45
Trp Gln Arg S r Trp Arg Tyr Trp His Asp Glu Gln Gly Met Ser Glu
          50           55           60

```

- 43 -

7. The protein according to claim 6, wherein said protein affects the structural organization or functional integrity of said mature virion by steric hindrance or enzymatic disturbance of said virion and/or by affecting viral protein interactions responsible for infectivity and/or viral replication.

8. The protein according to claim 1, which comprises a sufficient number of amino acids of an amino acid sequence, a functional derivative or fragment thereof, selected from the group consisting of:

Met Glu Gln Ala Pro Glu Asp Gln Gly Pro Gln Arg Glu Pro His Asn  
1 5 10 15

Glu Trp Thr Leu Glu Leu Leu Glu Glu Leu Lys Asn Glu Ala Val Arg  
20 25 30

His Phe Pro Arg Ile Trp Leu His Gly Leu Gly Gln His Ile Tyr Glu  
35 40 45

Thr Tyr Gly Asp Thr Trp Ala Gly Val Glu Ala Ile Ile Arg Ile Leu  
50 55 60

Gln Gln Leu Leu Phe Ile His Phe Arg Ile Gly Cys Arg His Ser Arg  
65 70 75 80

Ile Gly Val Thr Gln Gln Arg Arg Ala Arg Asn Gly Ala Ser Arg Ser  
85 90 95

(SEQ ID NO:1);

Met Ala Glu Ala Pro Thr Glu Leu Pro Pro Val Asp Gly Thr Pro Leu  
1 5 10 15

Arg Glu Pro Gly Asp Glu Trp Ile Ile Glu Ile Leu Arg Glu Ile Lys  
20 25 30

Glu Glu Ala Leu Lys His Phe Asp Pro Arg Leu Leu Ile Ala Leu Gly  
35 40 45

Lys Tyr Ile Tyr Thr Arg His Gly Asp Thr Leu Glu Gly Ala Arg Glu  
50 55 60

Leu Ile Lys Val Leu Gln Arg Ala Leu Phe Thr His Phe Arg Ala Gly  
65 70 75 80

Cys Gly His Ser Arg Ile Gly Gln Thr Arg Gly Gly Asn Pro Leu Ser  
85 90 95

- 42 -

**CLAIMS:**

1. A protein for targeting into a mature HIV-1 and/or HIV-2 virion, comprising a sufficient number of amino acids of a Vpr protein, a Vpx protein, functional derivatives or fragments thereof, wherein said protein interacts with Gag-precursor protein of said mature virion; thereby being incorporated by said mature virion.

2. The protein according to claim 1, wherein said Gag-precursor protein the p6 protein domain.

3. The protein according to claim 1, which further comprises a molecule attached to said protein, functional derivative or fragment thereof to form a chimeric protein, wherein said chimeric protein is incorporated by said mature virion.

4. The protein according to claim 3, wherein ~~said molecule is a protein fragment covalently~~ attached to the N- or C-terminal of said protein.

5. The protein according to claim 4 wherein said protein fragment comprises an amino acid sequence effective in reducing HIV infectivity and/or replication.

6. The protein according to claim 5, wherein said protein fragment comprises an amino acid sequence having a RNase or protease activity, an amino acid sequence creating steric hindrance during virion assembly and morphogenesis, and/or affecting viral protein interactions responsible for infectivity and/or viral replication.

- 45 -

His Phe Pro Arg Ile Trp Leu His Gly Leu Gly Gln His Ile Tyr Glu  
           35                          40                          45

Thr Tyr Gly Asp Thr Trp Ala Gly Val Glu Ala Ile Ile Arg Ile Leu  
           50                          55                          60

Gln Gln Leu Leu Phe Ile His Phe  
   65                          70                          (SEQ ID NO:4);  
 or

a sufficient number of amino acids of the following  
 amino acid sequence, a functional derivative or  
 fragment thereof:

Leu Gln Arg Ser Pro Glu Pro Thr Ala Pro Pro Glu Glu Ser Phe Arg  
   1                          5                          10                          15

Ser Gly Val Glu Thr Thr Thr Pro Pro Gln Lys Gln Glu Pro Ile Asp  
           20                          25                          30

Lys Glu Leu Tyr Pro Leu Thr Ser Leu Arg Ser Leu Phe Gly Asn Asp  
           35                          40                          45

Pro Ser Ser Gln  
           50                          (SEQ ID NO:5).

12. A method for substantially reducing HIV  
 expression or replication in a patient infected with  
 HIV-1 or HIV-2, which comprises administration of at  
 least one therapeutic agent selected from the group  
 consisting of the protein according to claim 1 and DNA  
 sequences encoding said protein, to said patient in  
 association with a pharmaceutically acceptable  
 carrier.

13. The method according to claim 12, wherein  
 said administration of a DNA construct harboring a  
 Vpr/Vpx-chimeric protein or functional derivative  
 thereof or of a Vpr/Vpx-chimeric protein or functional  
 derivative thereof is effected via the blood stream.

14. The method according to claim 12, wherein  
 said administration is effected intracellularly

- 44 -

Ala Ile Pro Thr Pro Arg Asn Met Gln  
 100 105 (SEQ ID NO:2); and

Met Thr Asp Pro Arg Glu Thr Val Pro Pro Gly Asn Ser Gly Glu Glu  
 1 5 10 15

Thr Ile Gly Glu Ala Phe Ala Trp Leu Asn Arg Thr Val Glu Ala Ile  
 20 25 30

Asn Arg Glu Ala Val Asn His Leu Pro Arg Glu Leu Ile Phe Gln Val  
 35 40 45

Trp Gln Arg Ser Trp Arg Tyr Trp His Asp Glu Gln Gly Met Ser Glu  
 50 55 60

Ser Tyr Thr Lys Tyr Arg Tyr Leu Cys Ile Ile Gln Lys Ala Val Tyr  
 65 70 75 80

Met His Val Arg Lys Gly Cys Thr Cys Leu Gly Arg Gly His Gly Pro  
 85 90 95

Gly Gly Trp Arg Pro Gly Pro Pro Pro Pro Pro Pro Gly Leu Val  
 100 105 110

(SEQ ID NO:3).

9. The protein according to claim 8, wherein said functional fragment thereof comprises a sequence of about 25 to about 63 amino acids.

10. A protein for substantially preventing replication of mature HIV-1 and HIV-2 virions, which comprises a Vpr/Vpx protein fragment, p6 protein, a p6 protein fragment, or functional derivatives thereof, wherein said protein interferes with the incorporation of native Vpr/Vpx into HIV-1 and/or HIV-2 virions by interacting with native Gag-precursor or with native Vpr or Vpx protein.

11. The protein according to claim 10, which comprises a fragment of the amino acid sequence or a functional derivative thereof:

Met Glu Gln Ala Pro Glu Asp Gln Gly Pro Gln Arg Glu Pro His Asn  
 1 5 10 15

Glu Trp Thr Leu Glu Leu Leu Glu Glu Leu Lys Asn Glu Ala Val Arg  
 20 25 30



- 47 -

20. A pharmaceutical composition for reducing HIV expression in a patient infected with HIV-1 or HIV-2, which comprises an effective amount of said protein according to claim 1 in association with a pharmaceutically acceptable carrier.

21. A pharmaceutical composition for reducing HIV expression in a patient infected with HIV-1 or HIV-2, which comprises a therapeutic amount of said protein according to claim 10, in association with a pharmaceutically acceptable carrier.

22. A vector comprising:

(a) a DNA segment encoding a protein which interferes with the incorporation of native Vpr/Vpx into HIV-1 and/or HIV-2 virions, comprising a sufficient number of amino acids of a Vpr protein, a Vpx protein, a p6 protein, functional derivatives or fragments thereof; and

(b) a promoter upstream of said DNA segment.

---

23. The vector according to claim 22, wherein said DNA segment encodes a Vpr/Vpx-chimeric protein.

24. The vector of claim 23, wherein said vector is adapted for expression of said chimeric protein in prokaryotic cells.

25. The vector of claim 23, wherein said vector is adapted for expression of said chimeric protein in eukaryotic cells.

- 46 -

through DNA transfection or administration of said protein.

15. A method according to claim 12, wherein said administration is effected by DNA transfection of said patient's hematopoietic cells followed by readministration of said transfected cells in said patient.

16. A method for substantially reducing HIV expression or replication in a patient infected with HIV-1 or HIV-2, which comprises administration of at least one therapeutic agent selected from the group consisting of said protein according to claim 10 and DNA sequences encoding said protein, to said patient in association with a pharmaceutically acceptable carrier.

17. The method according to claim 15, wherein said administration of a DNA construct harboring a Vpr/Vpx-chimeric protein or functional derivative thereof or of a Vpr/Vpx-chimeric protein or functional derivative thereof is effected via the blood stream.

18. The method according to claim 15, wherein said administration is effected intracellularly through DNA transfection or administration of said protein.

19. A method according to claim 15, wherein said administration is effected by DNA transfection of said patient's hematopoietic cells followed by readministration of said transfected cells in said patient.

1/15




FIG. 1

2/15

## VPR from HIVLAI

Met	Glu	Gln	Ala	Pro	Glu	Asp	Gln	Gly	Pro	Gln	Arg	Glu	Pro	His	Asn
1				5					10					15	
Glu	Trp	Thr	Leu	Glu	Leu	Leu	Glu	Glu	Leu	Lys	Asn	Glu	Ala	Val	Arg
			20					25					30		
His	Phe	Pro	Arg	Ile	Trp	Leu	His	Gly	Leu	Gly	Gln	His	Ile	Tyr	Glu
		35					40					45			
Thr	Tyr	Gly	Asp	Thr	Trp	Ala	Gly	Val	Glu	Ala	Ile	Ile	Arg	Ile	Leu
	50					55					60				
Gln	Gln	Leu	Leu	Phe	Ile	His	Phe	Arg	Ile	Gly	Cys	Arg	His	Ser	Arg
65					70					75					80
Ile	Gly	Val	Thr	Gln	Gln	Arg	Arg	Ala	Arg	Asn	Gly	Ala	Ser	Arg	Ser
				85					90					95	

## VPR from HIV2ROD

Met	Ala	Glu	Ala	Pro	Thr	Glu	Leu	Pro	Pro	Val	Asp	Gly	Thr	Pro	Leu
1				5					10					15	
Arg	Glu	Pro	Gly	Asp	Glu	Trp	Ile	Ile	Glu	Ile	Leu	Arg	Glu	Ile	Lys
			20					25					30		
Glu	Glu	Ala	Leu	Lys	His	Phe	Asp	Pro	Arg	Leu	Leu	Ile	Ala	Leu	Gly
		35					40					45			
Lys	Tyr	Ile	Tyr	Thr	Arg	His	Gly	Asp	Thr	Leu	Glu	Gly	Ala	Arg	Glu
	50					55					60				
Leu	Ile	Lys	Val	Leu	Gln	Arg	Ala	Leu	Phe	Thr	His	Phe	Arg	Ala	Gly
65					70					75					80

Cys	Gly	His	Ser	Arg	Ile	Gly	Gln	Thr	Arg	Gly	Gly	Asn	Pro	Leu	Ser
				85					90					95	
Ala	Ile	Pro	Thr	Pro	Arg	Asn	Met	Gln							
			100					105							

## VPX from HIV2ROD

Met	Thr	Asp	Pro	Arg	Glu	Thr	Val	Pro	Pro	Gly	Asn	Ser	Gly	Glu	Glu
1				5					10					15	
Thr	Ile	Gly	Glu	Ala	Phe	Ala	Trp	Leu	Asn	Arg	Thr	Val	Glu	Ala	Ile
			20					25					30		
Asn	Arg	Glu	Ala	Val	Asn	His	Leu	Pro	Arg	Glu	Leu	Ile	Phe	Gln	Val
		35					40					45			
Trp	Gln	Arg	Ser	Trp	Arg	Tyr	Trp	His	Asp	Glu	Gln	Gly	Met	Ser	Glu
	50					55					60				
Ser	Tyr	Thr	Lys	Tyr	Arg	Tyr	Leu	Cys	Ile	Ile	Gln	Lys	Ala	Val	Tyr
65					70					75					80
Met	His	Val	Arg	Lys	Gly	Cys	Thr	Cys	Leu	Gly	Arg	Gly	His	Gly	Pro
				85					90					95	
Gly	Gly	Trp	Arg	Pro	Gly	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Gly	Leu	Val
			100					105					110		



3/15

MEQAPEDGQPGQREPHNEWTLEELLEELKNEAVRHFPRIVLHGLGQHIYETYGDTWAGVEATIRILQQLLFTHFRIGCRHSRTGVTRRRARRNGASRS

10 20 30 40 50 60 70 80 90

PREDICTED HELIX

HxBRUWT

HxBRUA30F

HxBRUH33I

HxBRUR73S

HxBRUG75N

HxBRUR76/79

HxBRUR77/79

HxBRUR72/78

HxBRUE12,13PG

HxBRUE25K

HxBRUEA29,30FK


HxBRUIL63,64KR

HxBRULL68,70RK

HxBRUSR,79,80ID


FIG. 3

4/15



5/15

A



FIS-4C

6/15




Fig. 4D

Fig. 4E



7/15




FIG - 5B






FIG - 5A

8/15

FIG. 6



9/15

FIG. 7A



	Vpr incorporation
LL P10,11/L	+
LI/STOP	-
SI7/STOP	-
Y36/STOP	-
I44/STOP	-
P49/STOP	+

FIG. 7B

10/15




11/15




**FIG. 9**

12/15




FILE - 10

14/15

~~FIG. 12~~

15/15





## INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/CA 95/00510

A. CLASSIFICATION F SUBJECT MATTER  
 IPC 6 C12N15/49 C07K14/16 C12N15/62 C12N15/57 A61K39/21  
 C12N15/86 C12N5/10 //C12N9/52,C12N9/62,C12N9/64,C12N7/01

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	JOURNAL OF VIROLOGY 69 (5). 1995. 2759-2764. ISSN: 0022-538X, May 1995 KONDO E ET AL 'The p6-gag domain of human immunodeficiency virus type 1 is sufficient for the incorporation of Vpr into heterologous viral particles.' see the whole document --- -/--	1-11,22, 29

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*A\* document member of the same patent family

Date of the actual completion of the international search

14 November 1995

Date of mailing of the international search report

14. 12. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+ 31-70) 340-3016

Authorized officer

Hix, R



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA95/00510

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 12-19  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 12-19 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

.onal Application No

PCT/CA 95/00510

C.(Continuation) D CUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	KEYSTONE SYMPOSIUM ON GENE THERAPY AND MOLECULAR MEDICINE, STEAMBOAT SPRINGS, COLORADO, USA, MARCH 26-APRIL 1, 1995. JOURNAL OF CELLULAR BIOCHEMISTRY SUPPLEMENT 0 (21A). 1995. 395. ISSN: 0733-1959, KAPPES J C ET AL 'Targeting Foreign Proteins to HIV Particles Via Fusion with Vpr and Vpx.' see the whole document ---	1-29
P,X	JOURNAL OF VIROLOGY 68 (10). 1994. 6161-6169. ISSN: 0022-538X, October 1994 WU X ET AL 'Localization of the Vpx packaging signal within the C terminus of the human immunodeficiency virus type 2 gag precursor protein.' see the whole document ---	1,3-11, 22-29
X	JOURNAL OF VIROLOGY 67 (12). 1993. 7229-7237. ISSN: 0022-538X, December 1993 PAXTON W ET AL 'Incorporation of Vpr into human immunodeficiency virus type 1 virions: Requirement for the p6 region of gag and mutational analysis.' see the whole document ---	1-5, 7-11,22, 29
Y	see the whole document ---	6,23-28
X	VIROLOGY, vol. 199, no. 2, March 1994 pages 453-457, R.HORTON ET AL. 'HIV-2 viral protein X association with the gag p27 capsid protein' see the whole document ---	1,10,11, 22,29
X	WO,A,93 25235 (HARVARD COLLEGE) 23 December 1993	1,3-5, 7-11,20, 22,29
Y	see page 8, last paragraph - page 9, paragraph 1 ---	6,23-28
X,P	WO,A,95 16705 (UNIV PENNSYLVANIA ;WEINER DAVID B (US); LEVY DAVID NATHAN (US); RE) 22 June 1995 see the whole document ---	1,10,11, 20,22,29
X	WO,A,94 19456 (WEINER DAVID B ;LEVY DAVID NATHAN (US); REFAELI YOSEF (US)) 1 September 1994 see the whole document -----	1,10,11, 20,22